



MedicineWorld.Org

Your gateway to the world of medicine

Medicineworld.org: Where am I?

Home

News

Cancer News

About Us

Cancer

Health Professionals

Patients and public

Contact Us

Disclaimer

[Back to neurology news](#)

[Blogs list](#)

[Cancer blog](#)

Subscribe To Neurology News RSS Feed [RSS](#) [FEED](#) [?](#)

Where am I?

[Ads by Google](#)

[Psychology Quiz](#)

[Research Interest](#)

[Psychology Test](#)

[Psychology Studies](#)

[Market Research](#)



We've all experienced the feeling of not knowing where we are. Being disoriented is not pleasant, and it can even be scary, but luckily for most of us, this sensation is temporary. The brain employs many tricks to reorient us, keeping our confusion to a minimum and quickly pointing us in the right direction. Research has suggested that animals and young children mainly rely on geometric cues (e.g. lengths, distances, angles) to help them get reoriented. Human adults, however, can also make use of feature cues (e.g. color, texture, landmarks) in their surrounding area. But which method do we use more often? Psychology experts Kristin R. Ratliff from the University of Chicago and Nora S. Newcombe from Temple University conducted a set of experiments investigating if human adults have a preference for using geometric or feature cues to become reoriented.

[Psychology School Online](#)

Financial Planning is Available to All Students. Learn More Today!


online.argosy.edu

[PhD in Psychology](#)

Earn Your PhD in Psychology Online at Capella University. Learn More.

www.Capella.edu

Ads by Google



The first experiment took place in either a large or small white, rectangular room with a landmark (a big piece of colorful fabric) hanging on one wall. The study volunteers saw the researcher place a set of keys in a box in one of the corners. The volunteers were blindfolded and spun around, to become disoriented. After removing the blindfold, they had to point to the corner where the keys were. After a break, the volunteers were told the experiment would be repeated, even though they wouldn't watch the researcher hide the keys. Unbeknownst to them, during the break the scientists moved the landmark to an adjacent wall. This change forced the volunteers to use either geometric cues or feature cues, but not both, to reorient themselves and locate the keys. For the second experiment, the scientists used a similar method, except they switched room sizes (the volunteers moved from a larger room to a smaller room and vice versa)

during the break.

The results, reported in *Psychological Science*, a journal of the Association for Psychological Science, reveal that the brain does not have a distinct preference for certain cues during reorientation. In the first experiment, volunteers reoriented themselves by using geometric cues in the smaller room but used feature cues in the larger room. However, the volunteers who went from the larger room to the smaller room in the second experiment also relied on feature cues, searching for the landmark to become reoriented.

During the second experiment, the scientists surmise, the volunteers had a positive experience using feature cues in the large room, so they kept on relying on the landmark in the smaller room to become reoriented. These findings indicate that the brain takes into account many factors, including the environment and our past experiences, while determining the best way to reorient us to our surroundings.

Posted by: Daniel [Source](#)

[Psychology School Online](#)

Financial Planning is Available to All Students. Learn More Today!
online.argosy.edu

[PhD in Psychology](#)

Earn Your PhD in Psychology Online at Capella University. Learn More.
www.Capella.edu

Ads by Google

Did you know?

We've all experienced the feeling of not knowing where we are. Being disoriented is not pleasant, and it can even be scary, but luckily for most of us, this sensation is temporary. The brain employs many tricks to reorient us, keeping our confusion to a minimum and quickly pointing us in the right direction. Research has suggested that animals and young children mainly rely on geometric cues (e.g. lengths, distances, angles) to help them get reoriented. Human adults, however, can also make use of feature cues (e.g. color, texture, landmarks) in their surrounding area. But which method do we use more often? Psychology experts Kristin R. Ratliff from the University of Chicago and Nora S. Newcombe from Temple University conducted a set of experiments investigating if human adults have a preference for using geometric or feature cues to become reoriented.

Medicineworld.org: Where am I?

[Acute bacterial meningitis](#) | [Alzheimer's disease](#) | [Carpal tunnel syndrome](#) | [Cerebral aneurysms](#) | [Cerebral palsy](#) | [Chronic fatigue syndrome](#) | [Cluster headache](#) | [Dementia](#) | [Epilepsy seizure disorders](#) | [Febrile seizures](#) | [Guillain barre syndrome](#) | [Head injury](#) | [Hydrocephalus](#) | [Neurology](#) | [Insomnia](#) | [Low backache](#) | [Mental retardation](#) | [Migraine headaches](#) | [Multiple sclerosis](#) | [Myasthenia gravis](#) | [Neurological manifestations of aids](#) | [Parkinsonism](#) | [parkinson's disease](#) | [Personality disorders](#) | [Sleep disorders insomnia](#) | [Syncope](#) | [Trigeminal neuralgia](#) | [Vertigo](#) |

Copyright statement

The contents of this web page are protected. Legal action may follow for reproduction of materials without permission.

